

# Dangers of smoking whilst using oxygen therapy.

Important Information.



HIC/UKS/503895/BAAC/1112

**BOC Healthcare**

Patient Service Centre, Priestley Road, Worsley, Manchester, M28 2UT, United Kingdom  
Tel 0800 022 4332, [www.bochealthcare.co.uk](http://www.bochealthcare.co.uk)

The stripe symbol and the letters BOC are registered trade marks of The BOC Group Limited. Both BOC Limited and The BOC Group Limited are members of The Linde Group, the parent company of which is Linde AG. Reproduction without permission is strictly prohibited. © BOC Limited 2012

**BOC: Living healthcare**

Smoking around oxygen is extremely dangerous and may cause clothing and hair to catch fire and burn much more vigorously than in air. Never smoke or allow someone else to smoke nearby whilst using your oxygen equipment.



The picture above shows facial burns on a patient who lit a cigarette whilst using oxygen therapy.

### Oxygen enrichment

Your clothing could become enriched with oxygen as you use your medical oxygen. Clothing and materials which become enriched with oxygen will burn vigorously if ignited. Your clothing will continue to be enriched, even after you have turned off your oxygen supply.

Make sure you ventilate your clothing in the open air for at least 15 minutes before smoking or going near an open flame or source of ignition.

### Oils and grease

Oils, grease and materials contaminated with these substances are particularly hazardous in the presence of an oxygen enriched atmosphere, as they can ignite extremely easily and burn very violently.

- Ensure that your hands are clean and do not use oil-based creams and moisturisers when using your medical oxygen system This includes the use of oil-based lip moisturisers and make up.
- If using alcohol gel, please ensure that it is massaged in well, particularly between the webbing of fingers and allow time for evaporation to take place before handling oxygen equipment.
- Do not use oils or grease with your oxygen equipment.
- Use water-based products such as RoEezit\* or KY Jelly.

### Other hazards

- Do not use aerosols such as hair sprays or deodorants whilst using your oxygen equipment.

### Free NHS smoking helpline

Call to speak to a trained adviser for free on 0800 022 4332.

Lines are open Monday-Friday from 9am to 8pm; Saturday & Sunday from 11am to 5pm.

or visit their website [www.smokefree.nhs.uk](http://www.smokefree.nhs.uk)

Alternatively, for advice and support regarding smoking cessation, talk to your GP or nurse.

For more information on general fire safety, please visit the Fire Kills website at [www.direct.gov.uk/firekills](http://www.direct.gov.uk/firekills)

BOC supports the Fire Kills campaign.



Many thanks to the family of the patient pictured for their consent to use the photograph. \*RoEezit may need to be specially ordered from a pharmacy. All the safety information contained in this document was obtained from European Industrial Gases Association (EIGA)