How to administer Espranor[1]

Espranor is sensitive to moisture.
Make sure your hands are dry before handling the wafer.



1. Tear a square off the blister pack along the perforated lines.



3. Remove the wafer carefully from the foil and take out from the packaging immediately.



2. The foil is easily peelable. Do not force the wafer through the foil as it is fragile and can easily break. Instead, fold back the foil and then peel it off.



4. Place the wafer **on the tongue** and close your mouth. Allow it to remain there for a few seconds until it has dissolved. Try to avoid swallowing during the first 2 minutes.
Do not eat or drink for at least 5 minutes.

Job Bag Number: ADD/12/2017/178
Date of Prep: December 2017

MARTINDALE PHARMA®
Making lives better

Espranor®

(buprenorphine) oral lyophilisate

This leaflet is intended for patients that have been prescribed Espranor

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard.

By reporting side effects you can help provide more information on the safety of this medicine

Espranor is not interchangeable with other oral buprenorphine products and the dose of Espranor may differ from the dose of other buprenorphine products.

Job Bag Number: ADD/12/2017/178
Date of Prep: December 2017

MARTINDALE PHARMA®
Making lives better