



Pharmacy First for help with everyday health conditions

Pharmacy First Service for Acute Sore Throat due to Streptococcal infection in patients aged 5 years and over and adults

Service Specification and Guidance V2

Version	Details of changes	Date
2.0	<ul style="list-style-type: none"> All pharmacists including Independent Prescribing Pharmacists must provide the service in line with the service PGDs. The section dealing with possible complications, exclusion criteria and red flag symptoms has been removed. Pharmacists are directed to the service PGDs To operate under the new PGDs: <ul style="list-style-type: none"> Training now includes; NICPLD Sepsis e-learning module and NICPLD Safeguarding Advice & Guidance for Pharmacists recorded webinar ECHO training session recorded in November 2024 NOTE this training should be viewed alongside the most up-to-date versions of the PGDs (May 2026) <p>Key changes since recording:</p> <ul style="list-style-type: none"> Duration of all antibiotic courses is now FIVE days Pregnancy / suspected pregnancy under 16 years of age now excluded Breast feeding is no longer an exclusion for clarithromycin 	April 2026

Contents

1. Background.....	3
2. Service aims and objectives.....	4
3. Service description.....	4
4. Service outline	5
4.1 Patient Eligibility for the service	5
4.2 Pharmacy Eligibility for service.....	5
4.3 Pharmacist Eligibility to provide the service	6
4.4 Patient Consent	6
4.5 Accessing the service	6
4.6 Pharmacy First Consultation	6
(a) Assessment.....	7
(b) Rapid Antigen Detection Tests (RADT) / Consumables	10
(c) Possible Complications	10
(d) Treatment.....	12
(e) Advice for Patients	12
5. Service Formulary.....	13
6. Supply of medicine.....	15
7. Pharmacy First Consultation records	16
8. Premises.....	17
9. Professional responsibility.....	17
10. Training	18
11. Remuneration and Reimbursement	19
12. Service monitoring and post payment verification.....	20
13. Promotion of the service.....	20
14. Other terms and conditions.....	20
15. References.....	21
16. Appendices	21

1. Background

Acute sore throat is a common presentation in primary care and a leading cause of antibiotic prescribing. In the UK, almost 10% of patients consult their GP with sore throat every year and 8% of all acute prescribing relates to sore throat management. **Most acute sore throats are caused by a virus and result in mild self-limiting illness that can be self-managed with symptomatic treatment and do not require antibiotics.**

Bacterial sore throats are mainly caused by group A *Streptococcus* (GAS) and account for 5%–30% of all sore throat presentations.

The Department of Health's [Changing the Culture 2019 -2024 One Health strategy](#) for tackling antimicrobial resistance in NI includes measures to optimise the use of antimicrobials. The target set is a 15% reduction in antimicrobial use in humans by 2023/24. One key action to achieve this is to raise public awareness to encourage self-care and reduce expectations of antibiotics.

Sore Throat PILOT service 23/24

During the winter months November 2023 to March 2024 forty-three community pharmacies across NI provided a sore throat service to patients from the age of 5 years and above, who presented in the pharmacy with symptoms of sore throat.

A total of 6,768 consultations were undertaken as part of this pilot. In total 1,721 patients (25%) received treatment with antibiotics. A MOIC evaluation of the pilot service is available to view at this [link](#).

Identifying people with GAS-related acute sore throat (who may benefit from antibiotics) from clinical history and examination is difficult due to overlap of clinical signs and symptoms between bacterial and viral infections. It is generally accepted that clinical scoring criteria such as FeverPain score can help identify those with greater likelihood of GAS infection, but it has also been argued that scoring systems cannot replace clinical reasoning and judgement. To further support diagnosis of GAS-related pharyngitis, throat culture or rapid antigen detection testing (RADT) can be used.

A Pharmacy First Service is a service whereby patients are encouraged to consult with a participating community pharmacy rather than their GP for a defined list of common conditions. The pharmacist will give advice and (if appropriate) supply medication from an agreed formulary or refer the patient to another healthcare setting if necessary. Medicines, when

deemed necessary, are supplied free of charge.

This Pharmacy First Service for sore throat will facilitate the assessment and treatment of patients aged 5 years and over in the community pharmacy in line with [NI antimicrobial guidelines](#) and [NICE NG84 Sore throat \(acute\): antimicrobial prescribing](#).

2. Service aims and objectives

The aims of the service are to:

- Provide a more accessible, efficient and high-quality clinical pathway for patients with a sore throat.
- Better use pharmacist skills and free up GP time for more complex and urgent medical issues.
- Use point of care (POC) testing for group A *Streptococcus* (GAS) to guide management of the condition and potentially reduce unnecessary antibiotic prescribing.

The objectives of the service are to:

- Provide a timely and appropriate service for patients in the treatment of their condition and to identify patients who need onward referral to another healthcare professional.
- Provide a service which is acceptable to patients and community pharmacists and which is supported by GP practices.
- To assess how the use of diagnostic POC testing impacts the use of antimicrobials for the treatment of sore throat.
- Support the cost-effective use of medicines and health service resources in Primary Care in line with the NI formulary
- Promote the role of the community pharmacist as the first port of call for the management of sore throat in patients aged 5 years and over.

3. Service description

- The Pharmacy First Service allows eligible patients to use participating community pharmacies as the first port of call for the management of sore throat. The pharmacist advises, treats or refers patients according to their needs.
- The Pharmacy First Service is available to all eligible patients aged 5 years and over, registered with a GP in Northern Ireland, with the exception of temporary residents

and care home residents.

- Each Pharmacy First consultation must be carried out by an appropriately trained pharmacist with the patient directly. It should be face-to-face between the pharmacist and patient.
- The consultation must take place in a suitable private consultation area in the pharmacy (see section 8 'premises').

4. Service outline

4.1 Patient Eligibility for the service

The following persons are **eligible** for the service;

- Patients aged 5 years and over who are registered with a GP in Northern Ireland

The following persons are **not eligible** for the service;

- Temporary residents
- Patients in Care Homes (Nursing or Residential)

4.2 Pharmacy Eligibility for service

The service can only be provided from participating community pharmacies where the contractor:

- Holds a contract with the SPPG to deliver the service.
- Ensures that staff are trained, competent and available to deliver the service. Pharmacists must undertake relevant training to ensure clinical care competency prior to commencing service delivery (see section 10 'training').
- Ensures a Standard Operating Procedure (SOP) is in place to support delivery of the service in line with the service specification and guidance.
- Ensures Patient Group Directions (PGDs) for medicines relating to service delivery are organisationally authorised and signed by an appropriate authorising person.
- Ensures that the most up to date versions of service PGDs are used at all times. SPPG PGDs are available on the [BSO website](#).
- Ensures that the service is available during all of the pharmacy's opening hours, where practically possible.

- If in exceptional circumstances the responsible pharmacist on the day is not trained to offer the service, this should be communicated to the local GP practices and OOH medical centers to reduce inappropriate referral of patients to the pharmacy when the service is temporarily unavailable.

4.3 Pharmacist Eligibility to provide the service

This service can only be provided by pharmacists who are:

- Registered with the Pharmaceutical Society of Northern Ireland (PSNI). Pharmacist Independent Prescribers must be registered with the PSNI as an independent prescriber.
- Working in a pharmacy contracted to provide the service.
- Competent to provide the service (see section 10 'training').
- **IP pharmacists are not required to sign the service PGDs; however, it is recommended that they refer to them in the delivery of the service.**

4.4 Patient Consent

- Before the consultation the pharmacist must provide patients with sufficient information to inform consent to avail of the service.
- The service privacy notice should be used to explain to the patient how their personal data will be used and shared. A copy should be supplied if requested.

4.5 Accessing the service

- Patients with symptoms seeking advice and /or treatment contact the pharmacy in person or by phone.
- The pharmacist arranges a consultation with the patient in person in the pharmacy.
- Patients may be referred into the service from their GP practice. Arrangements for this should be agreed in advance between the GP practice and the pharmacy. The pharmacist should contact the practice to give an outline of the service and share a copy of the GP flow chart (appendix 1). This and other training materials for GP practices are available on the BSO website.

4.6 Pharmacy First Consultation

Acute sore throat is a symptom of an underlying condition and should be accurately diagnosed

before considering treatment. It is self-limiting and often triggered by a viral infection of the upper respiratory tract. Symptoms can last for around a week, but most people will get better within this time without antibiotics, regardless of cause¹. Antibiotics for streptococcal sore throat decrease symptom duration by around 16 hours, however are indicated in some situations^{2,3}.

Sore throat is often associated with the common cold. It may also be a symptom of influenza or of infectious mononucleosis (glandular fever). Sore throat caused by glandular fever may take longer to resolve (usually within 1–2 weeks), with associated lethargy continuing for some time afterwards. The most common bacterial cause of sore throat is Group A beta- haemolytic Streptococcus (GAS). Non-infectious causes of sore throat are uncommon, and include physical irritation from gastro-oesophageal reflux disease or chronic cigarette smoke and hayfever².

In addition to soreness on swallowing, patients may experience:²⁻⁵

- a dry, scratchy throat
- bad breath
- swollen neck glands
- headache, malaise, rhinitis and cough
- nausea, vomiting and abdominal pain — may be present in children with pharyngitis, and people with tonsillitis
- hoarseness if there is laryngeal involvement
- fever (common in pharyngitis and tonsillitis)

(a) **Assessment** (see appendix 2 for summary flow chart)

Assessment of the person is required to ensure appropriate management. Differentiating a viral sore throat from that caused by GAS on the basis of examination is difficult.

The [FeverPain](#) criteria should be used along with examination of the person to determine the likelihood of streptococcal infection (and therefore the need for antibiotic treatment)². To further support diagnosis of GAS-related pharyngitis rapid antigen detection testing (RADT) can be used.

The [FeverPain](#) score is scored out of 5 depending on how many of the following are present:

1. fever in the last 24 hours
2. purulent tonsils

3. attend rapidly (patient attended within 3 days of the onset of symptoms)
4. severely inflamed tonsils
5. no cough or coryza (catarrhal inflammation of the mucous membrane in the nose)

The pharmacist should assess and score each patient using the FeverPain criteria and follow the guidance below:

FeverPain score of 0 or 1

- DO NOT offer a RADT
- Do not offer antibiotics²
- Refer to the 'Symptom Relief' and 'Advice for Patients' sections below
- Advise the person to return to the pharmacy if symptoms do not improve after 7 days (or earlier if symptoms worsen)
- Seek advice from GP, OOH or ED if the person becomes systemically very unwell².

FeverPain score of 2 or 3

Options:

- Consider if the patient is likely to benefit from antibiotic treatment and, where this is the case, carry out a RADT. **See below for options if RADT is positive or negative.**
- If the patient is less likely to benefit from antibiotics, refer to the 'Symptom Relief' and 'Advice for Patients' sections below.

FeverPain score of 2 or 3

with a POSITIVE RADT result for Strep A

Options:

- Watch and wait if practical:
 - advise the person to return to the pharmacy for reassessment if symptoms fail to improve over the next 48 hours
 - advise seeking advice from GP, OOH or ED if the person becomes systemically very unwell²
 - refer to the 'Symptom Relief' and the 'Advice for Patients' section below

- o Consider supplying antibiotic with advice, depending on clinical condition; bearing in mind other circumstances (e.g. weekend/bank holiday), the unlikely event of complications if antibiotics are not taken and possible adverse effects:² **When deciding whether or not to supply an antimicrobial, consider the risk of antimicrobial resistance for individual patients and the population as a whole.**
 - If an antibiotic is not supplied advise the person to return to the pharmacy for reassessment if symptoms fail to improve over the next 48 hours
 - advise seeking advice from GP, OOH or ED if the person becomes systemically very unwell²
 - refer to the 'Symptom Relief' and 'Advice for Patients' sections below.

FeverPain score of 4 or 5

- o Carry out a RADT

FeverPain score of 4 or 5

with a POSITIVE RADT result for Strep A

- o If the person is not systemically very unwell; not showing signs of a more serious condition; and not at high risk of complications:
 - consider supplying antibiotic with advice, depending on clinical condition; bearing in mind other circumstances (e.g. weekend/bank holiday), the unlikely event of complications if antibiotics are not taken and possible adverse effects.² **When deciding whether or not to supply an antimicrobial, consider the risk of antimicrobial resistance for individual patients and the population as a whole.**
 - advise seeking advice from GP, OOH or ED if the person becomes systemically very unwell²
 - refer to the 'Symptom Relief' and 'Advice for Patients' sections below.

- o If antibiotics are not supplied:
 - advise the person to return to the pharmacy if symptoms fail to improve over the next 48 hours for reassessment.
 - advise seeking advice from GP, OOH or ED if symptoms worsen rapidly or

significantly, or the person becomes systemically very unwell²

- refer to the 'Symptom Relief' and 'Advice for Patients' sections below.
- If the person is systemically very unwell or showing signs of a more serious condition or at high risk of complications refer immediately as appropriate.

FeverPain score of 2, 3, 4 or 5

Options if RADT test is negative

- Reassure the patient that the sore throat is not likely to be bacterial and therefore antibiotics will be unlikely to help.
- If there is worsening of symptoms or new symptoms or symptoms fail to improve over the next 48 hours, advise the person to return to the pharmacy for reassessment.
- Advise seeking advice from GP, OOH or ED if the person becomes systemically very unwell²
 - Refer to the 'Symptom Relief' and 'Advice for Patients' sections below.

(b) Rapid Antigen Detection Tests (RADT) / Consumables

- Where appropriate, the Pharmacist will offer a RADT to patients presenting with a sore throat whose FeverPain score indicates possible Streptococcus A infection as indicated in the 'assessment' section above.
- Suitable Rapid Antigen Detection Tests (RADT) must be available during the sore throat consultation.
- Pages 8 to 14 of the [NICE DG38 Rapid tests for group A streptococcal infections in people with a sore throat diagnostics guidance](#) outlines further information in relation to a variety of RADT tests available to purchase.
- Other consumables which may be required include PPE e.g. gloves and masks, tongue depressors and a thermometer. Please refer to [PHA Infection Control Manual](#)

(c) Possible Complications / exclusions / red flag symptoms

Full details of exclusion criteria, red flag symptoms and when to refer to another health professional **must** be accessed in service PGDs:

[Phenoxymethylpenicillin PGD](#)

[Clarithromycin PGD](#)

[Erythromycin PGD](#)

Action to be taken if a patient is excluded – explain the reasons for exclusion to the patient and record reasons for exclusion in the appropriate clinical record e.g. patients PMR or paper copy of a consultation form.

(d) Treatment

Symptom Relief

- Paracetamol and/or Ibuprofen (if suitable) can be supplied to help ease pain and fever (see formulary section 5)
- Medicated lozenges containing either a local anesthetic, NSAID or an antiseptic may help with pain in adults. These may be purchased in the pharmacy.
- There is little evidence for benzydamine gargles /spray^{2,3}
- There is no evidence for zinc lozenges, herbal remedies or acupuncture^{2,3,5}
- Adults and older children may find sucking hard sweets, ice cubes or ice lollies provide symptomatic relief⁴
- Adults can try a warm saline gargle (half a teaspoon of salt in a glassful of warm water) at frequent intervals, but do not swallow. This is not suitable for young children⁴

Supply of antibiotics

- Based on the Pharmacist's clinical assessment of the patient and the outcome of any RADT, they will determine whether it is appropriate to supply an antibiotic. See appendix 4 for Antibiotic Decision Pathway
- First line antibiotic is phenoxymethylpenicillin unless the patient has a true allergy.
 - Approximately 1 in 100 people have a true penicillin allergy. However, 1 in 10 people have either been told or have assumed they have a penicillin allergy. This means that about 9 in 10 persons 'labelled' with a penicillin allergy will not be allergic. For many people, their reaction happened many years ago but was never further investigated. The label of penicillin allergy has just continued throughout their life.
 - Side effects are not the same as allergies
 - Link to Royal Pharmaceutical Society penicillin allergy check list [Penicillin allergy checklist | RPS \(rpharms.com\)](#)
 - Link to a useful patient information leaflet <https://antibioticresearch.org.uk/wp->

content/uploads/2021/10/Penicillin-Delabelling-Patient-Leaflet_21Apr21.pdf

- Non-IP pharmacists will determine appropriate supply of an antibiotic within the terms of the service PGDs.
- IP pharmacists will use their clinical judgement in the diagnosis and supply of appropriate antibiotics and be guided by the terms of the service specification and relevant PGDs.
 - Where the supply of an antibiotic is not appropriate, pharmacists should consider supply of symptomatic treatment, paracetamol and/or ibuprofen if appropriate.

(e) Advice for Patients

All patients must be offered advice and [leaflets](#) related to antimicrobial stewardship, especially regarding viral versus bacterial infections, and the self-limiting nature of the symptoms, regardless of the need and/or results of RADT.

General advice

- Colds, most coughs, sinusitis, ear infections, sore throats, and other infections often get better without antibiotics, as your body can usually fight these infections on its own⁶.
- Sore throat usually gets better within 7 days, with or without antibiotics⁶.
- Taking antibiotics makes bacteria that live inside your body more resistant so the antibiotics may not work when you really need them⁶.
- Antibiotics can cause side effects such as rashes, thrush, stomach pains, diarrhoea, reactions to sunlight, other symptoms, or being sick⁶.
- Provide TARGET information leaflet: [Respiratory tract infection resource suite: Patient facing materials\(rcgp.org.uk\)](#)

Advice if antibiotics are NOT supplied

- Return to the pharmacy if symptoms do not improve after 7 days (48 hours if FeverPain score of 2 or more), or earlier if symptoms worsen.
- Seek advice from GP, OOH or ED if the person becomes systemically very unwell².

Advice if antibiotics ARE supplied:

- Seek advice from GP if symptoms worsen or do not improve within 3– 5 days; seek advice from GP, OOH or ED if the person becomes systemically very unwell².

Lifestyle advice

- Rest and take simple painkillers at regular intervals to relieve pain and fever^{4,5}
- Avoid smoking and smoky environments⁴
- If you have a high temperature or you do not feel well enough to do your normal activities, try to stay at home and avoid contact with other people until you feel better⁴
- Drink plenty of water to avoid dehydration⁴
- Eat cool and soft foods⁵
- Hot drinks should be avoided as these can exacerbate pain²
- Children may return to school or day care after fever has resolved and they are no longer feeling unwell, and/or after taking antibiotics for at least 24 hours²

5. Service Formulary

First line antibiotic:

Phenoxymethylpenicillin (suitable for all patients who **DO NOT** have a penicillin allergy, **including** those aged 16 or over who are pregnant/breastfeeding)

Medication	Phenoxymethylpenicillin		
Formulation	250 mg tablets	250 mg/5 mL oral solution*	125mg/5ml oral solution*
Legal class	POM	POM	POM
Quantity to supply	Appropriate quantity for 5 days (see PGD)	Appropriate quantity for 5 days (see PGD)	Appropriate quantity for 5 days (see PGD)
Dosing instructions	<ul style="list-style-type: none"> • 125mg four times a day in those aged 5 years • 250 mg four times daily in those aged 6 to 11 years • 500 mg four times daily in those aged 12 years and over 		
	Duration of treatment is 5 days		

*sugar-free formulations should be supplied where possible

See [PGD](#) for counselling advice for Phenoxymethylpenicillin

Second line antibiotics:

Option A: Clarithromycin (where phenoxymethylpenicillin is not appropriate due to hypersensitivity and the person is NOT pregnant)

Medication	Clarithromycin			
Formulation	250mg tablets	500 mg tablets	125 mg/5 mL oral suspension*	250 mg/5 mL oral suspension*
Legal class	POM	POM	POM	POM
Quantity to supply	20	10	Appropriate quantity for 5 days (see PGD)	Appropriate quantity for 5 days (see PGD)
Dosing instructions	12 years and over: 500 mg twice daily for 5 days	12 years and over: 500 mg twice daily for 5 days	Children 5 – 11 years	Dose
			Body weight:	
			Up to 8kg	7.5 mg/kg twice daily
			8 – 11kg	62.5 mg twice daily
			12 – 19kg	125 mg twice daily
			20 - 29kg	187.5 mg twice daily
30 – 40kg	250 mg twice daily			
Duration of treatment is 5 days				

*sugar-free formulations should be supplied where possible

Option B: Erythromycin (aged 16 years and over who are pregnant/suspected pregnant and where phenoxymethylpenicillin is not appropriate due to hypersensitivity)

Medication	Erythromycin			
Formulation	250 mg tablets	500 mg tablets	125 mg/5 mL oral suspension*	250 mg/5 mL oral suspension*
Legal Class	POM	POM	POM	POM
Quantity to Supply	40	20	Appropriate quantity for 5 days (see PGD)	Appropriate quantity for 5 days (see PGD)
Dosing Instructions	<ul style="list-style-type: none"> 16 years and over: 500 mg four times a day 			
	Duration of treatment is 5 days			

*sugar-free formulations should be supplied where possible
^including gastro resistant tablets

See PGDs for counselling advice for [Clarithromycin](#) and [Erythromycin](#)

Analgesics (Ibuprofen is not suitable for patients who are pregnant or breast feeding)

Analgesia	Paracetamol		
Generic name	Paracetamol 120mg/5mL S/F oral suspension	Paracetamol 250mg/5mL S/F oral suspension	Paracetamol 500 mg tablets
Legal class	P	P	P
Pack size	100 ml	200 ml	32 tablets
Dosing instructions and advice	As per pack	As per pack	As per pack

Analgesia	Ibuprofen		
Generic Name	Ibuprofen 100 mg/5ml S/F oral suspension	Ibuprofen 200 mg tablets	Ibuprofen 400 mg tablets
Legal Class	P	P	P
Pack Size	100 ml	24 tablets	24 tablets
Dosing instructions and advice	As per pack	As per pack	As per pack

6. Supply of medicine

- a. On occasion a pharmacist will decide that a patient's symptoms are such that a supply of medicine(s) is indicated. Where this is the case the medicine(s) should be selected from the agreed formulary. Where a patient expresses a preference for an OTC/Pharmacy product which is not included in the agreed formulary and the pharmacist considers that such a supply is appropriate, the pharmacist can sell that product to the patient and the consultation shall still be considered to be within the terms of service provided that a record of the consultation is made.
- b. Where a medicine is supplied it shall be appropriately labelled and the pharmacist shall counsel the patient regarding its safe and effective use.
- c. Pharmacists must ensure medicines supplied comply with current good practice guidelines e.g.: Pharmaceutical Society guidance available at <http://www.psn.org.uk/publications/code-of-ethics-and-standards/> MHRA Drug Safety Advice <https://www.gov.uk/drug-safety-update>
Pack/product updates & individual SPCs <https://www.medicines.org.uk/emc/>
- d. When treatment is required and appropriate it should be selected from the formulary and supplied in one of two ways:
 - > The Pharmacist Independent Prescriber writes a prescription for the

medicine which is dispensed in accordance with the relevant SOP.

- Medicines prescribed in this way may be dispensed from Prescription Only Medicine (POM) packs.

Pharmacist Independent Prescribers working in a community pharmacy can only prescribe medicines as part of Pharmacy First Services and only those medicines listed in the service formulary.

- The non-IP Pharmacist supplies the medicine in line with the service PGDs and completes a Pharmacy Voucher (PV). Supply is made in accordance with the relevant SOP.
 - POM antibiotics must be supplied in line with service PGDs and analgesics must be supplied in OTC / P packs in line with product licenses.

Orders for prescription pads and PV1s must be placed on line on the BSO website at <https://hscbusiness.hscni.net/services/2540.htm>

7. Pharmacy First Consultation records

- a. All Pharmacy First consultation records must be full, accurate and contemporaneous
- b. In cases where antibiotics have not been supplied on initial assessment and the patient returns to the pharmacy for reassessment section 7 of the consultation record must be completed.
- c. A record of the consultation must be retained in the pharmacy and be available to SPPG for monitoring and audit purposes. In all cases a copy of the consultation form must be transferred securely to the patient's GP; where practical within 24/48 hours. Local arrangements for the secure transfer of patient data should be in place.
- d. All records must be kept for the time periods in line with the DOH Good Management, [Good records guidelines](#)
 - Adults - 8 years after the conclusion of treatment
 - Children and young people – Until the patient's 25th birthday or 26th if the young person was 17 at the conclusion of treatment or 8 years after death.
- e. IP prescriptions should be coded with normal drug tariff codes and submitted along with the usual prescription bundle to BSO for payment.
- f. PVs should be coded with normal drug tariff codes and the Pharmacy First code

97003/1 should be added. These PVs should be processed in line with other pharmacy vouchers and sent to BSO for payment.

- g. A record of all consultations should be made on the monthly claim form which should be emailed to local integrated care offices for processing and payment of service fees. See appendix 6 for a copy of monthly claim form, which should be submitted in a timely manner, in the first week of the month following the consultation.

Contact Details for Local Integrated Care Offices:				
Belfast	South Eastern	Southern	Northern	Western
12-22 Linenhall Street Belfast BT2 8BS	12-22 Linenhall Street Belfast BT2 8BS	Tower Hill Armagh. BT61 9DR	County Hall 182 Galgorm Road Ballymena BT42 1QB	Gransha Park House 15 Gransha Park Clooney Road Londonderry BT47 6FN
Tel: 028 9536 3926	Tel: 028 9536 3926	Tel: 028 9536 2104	Tel: 028 9536 2812	Tel: 028 9536 1082
pharmacyservicesbelfast@hscni.net	pharmacyservicesse@hscni.net	pharmacyservicesouth@hscni.net	pharmacyservicesnorth@hscni.net	pharmacyserviceswest@hscni.net

8. Premises

Pharmacies participating in the Pharmacy First Service must have a consultation area suitable for carrying out throat examination and swab, and meets the following requirements:

- a. The consultation area should be where both the patient and pharmacist can sit down together.
- b. The patient and pharmacist should be able to talk at normal speaking volumes without being overheard by another person (including pharmacy staff).
- c. The consultation area should be clearly designated as an area for confidential consultations, distinct from the general public areas of the pharmacy.
- d. The consultation area must provide equal access to all patients who may wish to avail of the Pharmacy First Service.
- e. **All** Pharmacy First sore throat consultations must take place in the consultation area.
- f. The pharmacy contractor should ensure that infection prevention and control measures around cleaning and decontamination requirements as recommended by PHA are followed. Please refer to [PHA Infection Control Manual](#)

9. Professional responsibility

- a. It is the responsibility of individual pharmacists to have suitable indemnity insurance cover (see appendix for Pharmacist Independent Prescribers).
- b. At all times the pharmacist will be required to preserve patient confidentiality in line with their responsibilities as members of the Pharmaceutical Society of Northern

Ireland and GDPR regulations.

- c. Access to patient's confidential medical records on NIECR: Pharmacists should only access the records of patients who have consented to avail of the service. Please be aware for both user and patient safety and governance, NIECR is heavily audited.
- d. At no point does this service abrogate the professional responsibility of the individual pharmacist. They must use their professional judgement at all times.
- e. The responsible pharmacist on the day is responsible for ensuring that the service is delivered in line with the service specification and guidance.
- f. Any complaints relating to the service should be dealt with in line with the participating pharmacy's complaints SOP.
- g. Pharmacists acting in the dual role of prescribing and supplying medicines should follow the joint [RCN and RPS Guidance on Prescribing, Dispensing, Supplying and Administration of Medicines](#).

10. Training

All pharmacists must undertake training necessary to meet the competency required to provide the service. It is the responsibility of the individual to keep up-to-date with continued professional development and to work within the limitations of individual scope of practice.

Mandatory training requirements:

1. Training was provided by SPPG via ECHO on 5th November 2024. This was recorded and is available on the ECHO Moodle site for pharmacists unable to attend the live event. All pharmacists planning to provide this service must view this recording prior to service delivery.

NOTE this training should be viewed alongside the most up-to-date versions of the service PGDs (1st May 2026).

Key changes since recording:

- **Duration of all antibiotic courses is now FIVE days**
- **Pregnancy / suspected pregnancy under 16 years of age now excluded**
- **Breast feeding is no longer an exclusion for clarithromycin**

2. [NICPLD Sepsis e-learning](#)

3. NICPLD [Safeguarding Advice & Guidance for Pharmacists](#) recorded webinar

Other recommended therapeutics training:

- The TARGET Antibiotics Toolkit training resources available at [TARGET tools to train prescribers: TARGET antibiotic toolkit training resources \(rcgp.org.uk\)](#) which includes a video presentation <https://youtu.be/wLFJtcn5S7g>
- NICE guidance on [antimicrobial stewardship](#), NICE Guideline, [NG84. Sore throat \(acute\) in adults: antimicrobial prescribing](#) and NICE [CKS sore throat - acute](#)
- NICPLD on line modules [Antimicrobial Stewardship](#)NIECR training:

NIECR training:

Training guides and FAQs on the use of NIECR are available at:

<https://bso.hscni.net/directorates/operations/family-practitioner-services/pharmacy/contractor-information/contractor-communications/hscb-services-and-guidance/northern-ireland-electronic-care-record-niecr/>

- An NIECR eLearning module is now also available via NICPLD - [NICPLD: Open learning](#) (Under Pharmacy Practice)

PGD training:

- [Patient Group Directions - elearning for healthcare \(e-lfh.org.uk\)](#)
Freely available to all without registration
- NICE MPG 2 - [Overview | Patient group directions | Guidance | NICE](#)

It is good practice for the pharmacy contractor to keep documentary evidence that pharmacy staff involved in the provision of the service are competent and remain up to date with regards to the specific skills and knowledge that are appropriate to their role, and to the aspects of the service they are delivering.

11. Remuneration and Reimbursement

The fees payable to pharmacy contractors for this service are:

- One off service set-up payment of £200 per pharmacy contractor
- A consultation fee of £32 per consultation. This fee includes the cost of RADTs and

any other necessary consumables.

- The cost of medicines supplied will be reimbursed on submission to BSO of the prescriptions or pharmacy vouchers.

12. Service monitoring and post payment verification

- a. The pharmacy contractor will be required to submit all records requested by SPPG in relation to the Pharmacy First Service within 14 days of receipt of the request.
- b. The pharmacy contractor is required to co-operate on a timely basis in respect of any review or investigation being undertaken by SPPG / BSO regarding the Pharmacy First Service.
- c. In the event where SPPG cannot assure claims relating to the provision of the Pharmacy First Service recovery of the payment will be sought.

13. Promotion of the service

SPPG will provide printed A3 and A4 posters for use within the pharmacy. Pharmacies may also wish to promote the service on Twitter and Instagram using the resources available on the PCI and BSO websites. The pharmacy contractor shall not publicise the availability of the service other than using any materials specifically provided by SPPG without the prior agreement of the SPPG or in any way which is inconsistent with the professional nature of the service.

14. Other terms and conditions

- a. The pharmacy contractor shall not give, promise or offer to any person any gift or reward as an inducement to or in consideration of his/her registration with the service.
- b. The pharmacy contractor shall not give, promise or offer to any person engaged or employed by him any gift or reward or set targets, against which that person will be measured, to recruit patients to the service
- c. The pharmacy contractor shall ensure that service provision is in accordance with professional standards.

15. References

1. National Institute for Health and Care Excellence. NICE Guideline, NG84. Sore throat (acute) in adults: antimicrobial prescribing. Jan 2018. Available at: <https://www.nice.org.uk/guidance/ng84>
2. NICE Clinical knowledge summaries. Sore throat - acute. Jan 2023. Available at: <https://cks.nice.org.uk/topics/sore-throat-acute/>.
3. Patient. Sore throat: causes, symptoms and treatment. May 2022. Available at: <https://patient.info/doctor/sore-throat-pro>.
4. NHS. Sore throat. Feb 2021. Available at: <https://www.nhs.uk/conditions/sore-throat/>.
5. NHS. Tonsillitis. Feb 2021. Available at: <https://www.nhs.uk/conditions/tonsillitis/>.
6. Royal College of General Practitioners. RTI leaflet. A leaflet for treating respiratory tract infections. Nov 2021. Available at: <https://elearning.rcgp.org.uk/mod/book/view.php?id=12647&chapterid=478>

16. Appendices

Appendices available on the BSO website [here](#)

Appendix 1 – GP Flowchart

Appendix 2 – Service summary flow chart

Appendix 3 – Antibiotic decision pathway

Appendix 4 - Consultation form

Appendix 5 – Monthly claim form

Appendix 6 – Independent Prescribers